

Neutrino and neutron spectroscopy using trapped ions

Dr. Nicholas Scielzo

Lawrence Livermore National Laboratory

Monday, May 5 ♦ 4 P.M.

Room 124 Nieuwland Science Hall

The neutrinos and neutrons emitted in nuclear beta decay can be precisely studied using radioactive ions held in a radiofrequency-quadrupole ion trap. When a radioactive ion decays in the trap, the recoil-daughter nucleus and emitted particles emerge from the $\sim 1\text{-mm}^3$ trap volume without scattering and propagate unobstructed through vacuum. This allows the momentum and energy of particles that would otherwise be difficult (or even impossible) to detect to be reconstructed from the momentum imparted to the recoiling nucleus. Measurements of beta-neutrino angular correlations can be made by taking advantage of the favorable properties of the ^8Li and ^8B beta decays and the benefits afforded by using trapped ions to allow an accurate determination of the direction and energy of each emitted neutrino. Beta-delayed neutron spectroscopy can be performed by circumventing the many difficulties associated with direct neutron detection and instead reconstructing the neutron emission probabilities and energy spectra from the time of flight of the recoiling nuclei. These novel techniques will have an important impact on improving our understanding of fundamental electroweak theory and the origin of the elements and will benefit applications of nuclear science such as nuclear energy and stockpile stewardship. Recent results for ^8Li and b-delayed neutron spectroscopy at CARIBU will be presented and future plans for these experiments will be discussed.

This work performed under the auspices of the U.S. Department of Energy by Lawrence Livermore National Laboratory under Contract DE-AC52-07NA27344.